



# Evaluation of anti-ulcerogenic activity in an Aqueous Extract obtained from *Bauhinia forficata* leaves

Giovanna Christina Costa Mazzeo<sup>1</sup>; Marcos Paulo de Oliveira Silva<sup>2</sup>; Luciana Lopes Guimarães<sup>1</sup>; Alba Regina Monteiro Souza Brito<sup>1,3</sup>; Walber Toma<sup>1,4\*</sup>

<sup>1</sup> Universidade Santa Cecília (UNISANTA), Programa de Pós-Graduação em Sustentabilidade de Ecossistemas Marinhos e Costeiros (PPG Ecomar), Santos, SP, Brasil.

<sup>2</sup> Universidade Santa Cecília (UNISANTA), Faculdade de Educação Física e Farmácia, Santos, SP, Brasil.

<sup>3</sup> Universidade Estadual de Campinas (UNICAMP), Laboratório de Produtos Naturais –Departamento de Biologia Funcional e Molecular –Campinas, SP, Brasil.

<sup>4</sup> Centro Universitário São Camilo (SÃO CAMILO), Curso de Farmácia – São Paulo, SP, Brasil.

## ABSTRACT

*Bauhinia forficata* Link, popularly known as pata-de-vaca, unha-de-vaca, casco-de-vaca, has been widely used in traditional medicine to treat several diseases. Leaves of *B. forficata* are used in popular medicine as a diuretic, hypoglycemic, tonic and cleanser, and to combat elephantiasis. However, despite the wide range of ethnopharmacological data surrounding the plant, there are no scientific data demonstrating a probable anti-ulcerogenic activity conferred by use of that species. The present study aimed to evaluate the antiulcer properties of an infusion of fresh leaves of *B. forficata* Link. From the leaves of *B. forficata*, an Aqueous extract (AqE) was obtained and the phytochemical analysis showed the presence of flavonols in this extract. In the gastric ulcer induced by administration of HCl-Ethanol model performed with four different doses of AqE (125, 250, 500 and 1000 mg.Kg<sup>-1</sup>), the AqE showed significant preventive activity (\*p<0.01) at doses of 1000 mg.Kg<sup>-1</sup>. The antiulcer activity of AqE (1000 mg.Kg<sup>-1</sup>) could also be demonstrated in experimental models of NSAID-bethanechol (\*\*p<0.001) and absolute ethanol (\*\*p<0.001). Moreover, AqE (1000 mg.Kg<sup>-1</sup>) promoted a significant increase (\*\*p<0.001) in the amount of gastric mucus. The data presented here demonstrated the potential gastroprotective activity from AqE, possibly attributed to the presence of flavonols in this extract. These results may serve as a support for the development of new treatments related to the pathology of gastric ulcer.

Keywords: Gastric ulcer. Cytoprotection. Flavonoids. *Bauhinia forficata*.

## INTRODUCTION

Peptic ulcer is a term used to describe a group of ulcerative disorders that occur in areas of the upper gastrointestinal tract that are exposed to acidic secretions and pepsin. This pathology is characterized by mucosal damage and is predominantly caused by *Helicobacter pylori*, antiplatelet agents such as acetylsalicylic acid, nonsteroidal anti-inflammatory drugs (NSAIDs) such as oral bisphosphonates, potassium chloride, immunosuppressive medications, serotonin reuptake inhibitors, alcohol consumption, and cigarette smoking (Golbabapour *et al.*, 2013).

Peptic ulcer represents a chronic health problem. Approximately 10% of the population have or will develop a peptic ulcer (Brito Júnior *et al.*, 2013). In Brazil, despite the absence of epidemiological records on ulcer cases, it is known that there are numerous cases involving this disease, meaning that it is a significant public health problem that has prompted a search for new substances with anti-ulcerogenic activity. Although a large arsenal of drugs with anti-ulcerogenic activity is already on the market, none results in 100% remission of ulcers with minimal side effects and without compromising the patient's wellbeing, which usually results in the chronic use of these drugs. Published studies have reported the widespread identification of new drugs derived from natural anti-ulcerogenic sources (Jamal *et al.*, 2006).

According to the World Health Organization, 80% of people in developing countries depend on traditional medicinal practices to meet and/or supplement their basic health needs (Pires *et al.*, 2011). Currently, despite marketing and encouragement from the pharmaceutical industry during the development of allopathic medicines, a large segment of the population in many countries continues to utilize complementary practices for their health care (Bonifácio *et al.*, 2014).

Brazil is an important reference point for the study of medicinal plants. With more than 56,000 species (excluding fungi), Brazil has one of the richest floras in the world – nearly 19% of the world's flora (Giulietti *et al.*,

*Autor correspondente:* Prof. Dr. Walber Toma, Universidade Santa Cecília (UNISANTA), Programa de Pós-Graduação em Sustentabilidade de Ecossistemas Marinhos e Costeiros (PPG Ecomar); Centro Universitário São Camilo (SÃO CAMILO), Curso de Farmácia. e-mail: walbertoma@unisanta.br, walbertoma@gmail.com

2005). Among this wide range of plant species, particularly relevant in the present context is *Bauhinia forficata* Link (Caesalpinioideae). This species corresponds to one of about 300 species of *Bauhinia*, and is also popularly known as Unha-de-Vaca, Pata-de-Vaca and Casco-de-Vaca. It is an ornamental tree because its showy white flowers are widely used in landscaping and urban forestry. Native to the Atlantic forest, this species is found from Rio de Janeiro to Rio Grande do Sul; therefore, its presence is very common on the coast of Santos city (Lorenzi, 2002).

*B. forficata* Link is a species that has been used in folk medicine in Africa, Asia and America for the treatment of various diseases (Pizzolatti, 2003). Moreover, an infusion of the leaves of *B. forficata* is used in popular medicine as a diuretic, hypoglycemic, tonic and cleanser, and to combat elephantiasis (Martins *et al.*, 1988). However, despite the wide range of ethnopharmacological data surrounding the plant, our literature review in databases of biomedical scientific information such as PubMed and Google Scholar showed that there are no data regarding a probable anti-ulcerogenic activity conferred by use of *B. forficata*.

Considering the need to search for new resources for the treatment of gastric ulcers and the exponential growth in scientific research about medicinal plants, the present study aimed to evaluate the antiulcer properties of an infusion of fresh leaves of *B. forficata* Link.

## MATERIALS AND METHODS

### *Plant Specimen and Extraction*

*B. forficata* Link leaves were obtained from Peruibe, São Paulo, Brazil (-24,267948, -46,959276 (latitude/longitude)) in March 2007 and were identified by Botanical Ms Paulo Salles Penteadó; a voucher specimen with number 4651 was deposited in the herbarium of the Universidade Santa Cecília (HUSC). The fresh leaves were submitted to an infusion extraction according to Pepato *et al.* (2002). In this process, the infusion with the leaves of this species was performed at a ratio of 150 g of fresh leaves/liter of water. Subsequently, the extracts were filtered and concentrated under vacuum to obtain an Aqueous Extract (AqE) from *B. forficata* leaves.

### *Phytochemical Analysis*

Tests for the presence of flavonoids in the AqE were carried out according to previously reported methods (Shinoda test, alkaline reagent test, ferric chloride test and oxalo-boric acid reaction) (Costa, 1982; Harborne, 1973; Harborne, 1998; Zuanazzi & Montanha, 2010).

### *Animals*

Male Wistar rats (150–250 g) and male Swiss mice (25–35 g), both obtained from the breeding facility of the Santa Cecília University (UNISANTA), were used. The animals were fed a certified Nuvilab® (Nuvital) diet with free access to tap water under standard conditions of 12 h dark–12 h light, humidity ( $60 \pm 1.0\%$ ) and temperature

( $21 \pm 1^\circ\text{C}$ ). Moreover, the animals were kept in cages with raised floors of wide mesh to prevent coprophagy. The protocols were approved by the Santa Cecília University Institutional Animal Care and Use Committee (CEUA – UNISANTA) under protocol number 53/07.

### *HCl/Ethanol-Induced Ulcer*

The anti-ulcerogenic activity of the AqE from *B. forficata* leaves was assessed in mice using this model as described by Mizui and Doteuchi (1983). Mice were divided into 3 groups which were fasted for 24 h prior to oral dosing with the vehicle, 0.9% Saline (10 mL.Kg<sup>-1</sup>), Lansoprazole (30 mg.kg<sup>-1</sup>), or AqE (125, 250, 500 and 1000 mg.Kg<sup>-1</sup>). Fifty minutes after the treatments, all animals received 0.2 ml orally of a 0.3 M HCl/60% EtOH solution. Animals were killed 1 h after the administration of HCl/EtOH solution; the stomachs were excised, inflated by an injection of saline (2 ml) and opened along the greater curvature. Then the stomachs were fixed in 5% formalin for 30 min and the ulcerative lesion index (ULI) was calculated by the methodology of Szelenyi and Thiemer (Szelenyi and Thiemer, 1978).

### *NSAID-Induced Gastric Ulcers in Cholinomimetic-Treated Mice*

The experiment was performed according to the method of Rainsford, 1978. In this model, gastric ulcer was induced using indomethacin (30 mg.kg<sup>-1</sup>, s.c.) and bethanechol (5 mg.kg<sup>-1</sup>, i.p.) administered to mice after a 24 h fast. AqE (1000 mg.kg<sup>-1</sup>), Lansoprazole (30 mg.kg<sup>-1</sup>) or 0.9% Saline (10 mL.Kg<sup>-1</sup>) were administered orally 30 min before the induction of a gastric ulcer. The animals were killed by cervical dislocation 4 h after treatment with the ulcerogenic agents; the stomachs were removed and inflated with 4% formalin in buffered saline, and the gastric damage was determined by the methodology of Szelenyi and Thiemer (1978).

### *Ethanol-induced gastric ulcer*

Rats were subjected to an ethanol-induced ulcer assay (Morimoto *et al.*, 1991). The rats were fasted for 24 h (free access to water) before the experiment. The animals were then randomly divided and received either oral AqE (1000 mg.Kg<sup>-1</sup>), Lansoprazole (30 mg.Kg<sup>-1</sup>) or 0.9% Saline (10 mL.Kg<sup>-1</sup>). After 30 minutes, all animals received 1 mL of absolute ethanol orally. One hour later, the animals were killed by cervical dislocation and their stomachs were removed to determine their ulcer index (Szelenyi & Thiemer, 1978).

### *Determination of Mucous in Gastric Contents*

This assay was performed according to the methodology described previously by Curtis *et al.* (1995), with some modifications. Mice were fasted for 24 h under anesthesia, with the abdomen incised and the pylorus ligated. Either AqE (1000 mg.kg<sup>-1</sup>), Carbenoxolone (200 mg.Kg<sup>-1</sup>) or vehicle (10 mL.Kg<sup>-1</sup>) were administered

intraduodenally after the pylorus ligation. The animals were killed by cervical dislocation 4 h after the drug treatments. The stomach contents were immersed in 10 mL of 0.02% Alcian Blue in 0.16 M sucrose/0.05 M sodium acetate, pH 5.8 and incubated for 24 h at 20°C. The Alcian Blue binding extract was centrifuged at 3000 rpm for 10 min. The absorbance of the supernatant was measured at 615 nm using a light spectrophotometer U/2000 (Hitachi, Japan). The free mucous in the gastric contents was calculated from the amount of Alcian Blue Binding (mg/wet tissue (g)).

#### Statistical Analysis

Results were expressed as the mean $\pm$  (S.E.M.) and statistical significance was determined by one-way analysis of variance (ANOVA) followed by Dunnett's post hoc test, with the minimum level of significance set at  $p < 0.05$ .

## RESULTS

Some studies have shown that *B. forficata* extract presented flavonoid compounds such as Kaempferol (a flavonol compound). The phytochemical screening of this study showed that AqE from *B. forficata* leaves presented flavonols in four different analyses (Table 1).

We can observe that the AqE from *B. forficata* leaves at doses of 1000 mg.Kg<sup>-1</sup> by the oral route prevented the gastric lesions induced by the administration of HCl Ethanol (\* $p < 0.01$ ), absolute ethanol (\*\* $p < 0.001$ ) and NSAIDS-Bethanecol (\*\* $p < 0.001$ ) (Table 2).

Still searching for a possible gastroprotective mechanism for the AqE from *B. forficata* leaves, we investigated the effects of this extract (1000 mg.Kg<sup>-1</sup>) on experimental protocols of mucus production. Pretreatment with AqE and Carbenoxolone (200 mg.kg<sup>-1</sup>) significantly (\*\* $p < 0.001$ ) increased the amount of adherent mucus in the gastric mucosa when compared to the control group (Figure 1).




Figure 1 - Effects of a vehicle, AqE, from *B. forficata* leaves and Carbenoxolone on adherent gastric mucus after pylorus-ligation in rats. ANOVA with Dunnett's test. \*\* $p < 0.001$  compared to the 0.9% Saline Control Group

## DISCUSSION

Flavonoids are some of the most common phenolics, widely distributed in plant tissues and often responsible, together with carotenoids and chlorophylls, for their blue, purple, yellow, orange and red colors. The flavonoid family

Table 1 - Phytochemical screening of Aqueous Extract (AqE) from *B. forficata* leaves

Phytochemical Analysis	Flavonoid Classification	Presence (+)
Shinoda	Flavonols	(+)
Oxalo-Boric	Flavonols	(+)
FeCl <sub>3</sub>	Flavonols	(+)
Alkaline Reagent Test	Flavonols	(+)

Table 2 - Effects of an aqueous extract (AqE) obtained from *B. forficata* Link leaves in different models of acute gastric lesion induced in rats or mice.

Gastric Ulcer Models	Treatment (p.o.)	n	Dose	ULI (mm)	Inhibition (%)
HCl-Ethanol in mice	0.9% Saline	07	10 mL.Kg <sup>-1</sup>	36.9 $\pm$ 11.0	---
	Lansoprazole	08	30 mg.Kg <sup>-1</sup>	12.7 $\pm$ 1.7**	65.6%
	AqE	07	125 mg.Kg <sup>-1</sup>	36.9 $\pm$ 3.8	---
	AqE	07	250 mg.Kg <sup>-1</sup>	30.0 $\pm$ 3.5	---
	AqE	07	500mg.Kg <sup>-1</sup>	29.7 $\pm$ 3.0	---
Ethanol in Rats	0.9% Saline	07	10 mL.Kg <sup>-1</sup>	94.6 $\pm$ 8.0	---
	Lansoprazole	07	30 mg.Kg <sup>-1</sup>	59.4 $\pm$ 7.4**	37.2%
	AqE	07	1000 mg.Kg <sup>-1</sup>	75.7 $\pm$ 6.1**	20%
NSAIDS Bethanecol in mice	0.9% Saline	08	10 mL.Kg <sup>-1</sup>	22.0 $\pm$ 2.1	---
	Lansoprazole	10	30 mg.Kg <sup>-1</sup>	11.0 $\pm$ 1.4**	50%
	AqE	08	1000 mg.Kg <sup>-1</sup>	16.0 $\pm$ 1.9**	27.3%

Data are presented as mean  $\pm$  S.D (n=7-10). ANOVA followed by Dunnett's test with \* $p < 0.01$ ; \*\*  $p < 0.001$

includes flavones, flavonols, iso-flavonols, anthocyanins, anthocyanidins, proanthocyanidins and catechins (Ferreira & Pinho, 2012; Rong, 2010).

These phenolic compounds (Flavonoids) belong to the recently popularized phytochemicals, chemicals derived from plant material with potentially beneficial effects on human health. The therapeutic effects of many traditional medicines may be related in many cases to the presence of these polyphenols (Manach *et al.*, 1997). For example, a wide variety of pharmacological activities have been reported for these substances, including antiviral (Critchfield *et al.*, 1996), anti-allergic (Cheong *et al.*, 1998), antiplatelet (Carotenuto *et al.*, 1997), anti-estrogenic, anti-carcinogenic, anti-inflammatory, anti-proliferative, antiangiogenic, and antioxidant properties, and their ingestion typically produces no or very little toxicity (Havsteen, 2002). Flavonoids are also reported to act in the gastrointestinal tract, having antispasmodic, anti-secretory, antidiarrheal and anti-ulcerative properties (Mota *et al.*, 2009). In our phytochemical screening, we found that AqE from *B. forficata* leaves presented flavonols in four different analyses (Table 1).

Experimental animals play an important role in the search for new drugs with protective properties. Considering that the etiology of ulcer is multifactorial, lesions in the gastric mucosa can be induced in different experimental

models through various mechanisms (Samonina, 2004). Some of the most widely used acute models for the evaluation of anti-ulcerogenic substances in animals are models of gastric lesion induced by HCl-ethanol, absolute ethanol and NSAIDs-bethanecol. In this work, we used these methodologies to evaluate the pharmacological activity of an AqE from *B. forficata* leaves.

Lee *et al.* (2010) related that the administration of HCl-ethanol produces ulcerative lesions and increases lipid peroxidation in the gastric mucosa, which plays a significant part in the pathogenesis of the mucosal lesions. Moreover, HCl-ethanol instillation induces severe epithelial desquamation, deep mucosal necrosis and submucosal edema associated with leucocyte accumulation. Brito Júnior *et al.* (2013) showed that the gastric lesions induced by absolute ethanol occur predominantly in the glandular portion of the stomach, as the result of a direct necrotizing action and, moreover, a lack of defense mechanisms such as secretion of bicarbonate and mucus and increased oxidative stress. The suppression of prostaglandin synthesis by nonsteroidal anti-inflammatory drugs (NSAIDs), such as indomethacin, is known to result in the increased susceptibility to mucosal injury and gastroduodenal ulceration. Indomethacin inhibits both cyclooxygenase<sup>1</sup> (COX<sup>1</sup>) and (COX-2) as well as the production of prostaglandins in the stomach and intestines to maintain the mucous lining of the gastrointestinal tract. Indomethacin, therefore, like other non-selective COX inhibitors, can cause peptic ulcers. Cholinomimetic agents (bethanechol) administered in association with NSAIDs have a synergistic effect on the gastric injury induced by increased secretion of acid and pepsin in the stomach (Lee *et al.*, 2010).

In our protocols, we observed that the AqE (1000 mg.Kg<sup>-1</sup>) by the oral route prevented gastric lesions induced by the administration of HCl Ethanol (\*p<0.01), absolute ethanol (\*\*p<0.001) and NSAIDs-Bethanecol (\*\*p<0.001) (Table 2).

Several works have shown that substances with antioxidant properties, such as flavonoids, may protect against the gastric-damaging effects caused by HCl-ethanol, absolute ethanol and NSAIDs-Bethanecol (Matsumoto *et al.*, 2004; Dekanski *et al.*, 2009). Hirano *et al.* (1994) demonstrated that a flavone protected the gastric mucosa of rats from injuries induced by ethanol. Moreover, Izzo *et al.* (1994) also obtained positive results in the treatment of gastric ulcer by utilizing flavonols. Most likely, the flavonols present in the AqE from *B. forficata* leaves are responsible for the anti-ulcerogenic activities evident in the experimental protocols demonstrated in this work.

In a review by Mota *et al.* (2009), the mechanisms of action of flavonoids such as flavonols in gastric protection were reported to include increased gastric blood flow and stimulated mucus synthesis in the gastric mucosa (\*\*p<0.001) (Figure 1). Most probably, the increase in adherent gastric mucus apparent after administration of the AqE from *B. forficata* leaves is possibly associated with the presence of flavonols in this extract. Considering the

lack of the cytoprotective treatments against gastric ulcers, new sources of cytoprotective treatments are vital. Thus, the data shown have an important bearing on new studies about probable cytoprotective treatments for gastric ulcers.

In conclusion, the AqE from *B. forficata* leaves showed preventive anti-ulcerogenic activity in mice in 3 ulcer models. Mucus secretion is involved in the gastroprotection exerted by this species, probably due to the flavonoids (flavonols) present in the plant. These results are important for the development of future cytoprotective treatments for gastric ulcers.

## ACKNOWLEDGMENTS

Toma W. acknowledges FAPESP (São Paulo Research Foundation) for the financial support (Grant. no. 2006/59888-7).

## CONFLICTS OF INTEREST

The authors of this article declare no conflict of interest regarding the data presented in the study.

## REFERENCES

- Bonifácio BV, Silva PB, Ramos MAS, Negri KMS, Bauab TM, Chorilli M. Nanotechnology-based drug delivery systems and herbal medicines: a review. *Int J Nanomed.* 2014;9:1–15.
- Brito Júnior FE, Oliveira DR, Bento EB, Leite LHI, Souza DO, Siebra AA, Sampaio RS, Martins AOPB, Ramos AGB, Tintino SR, Lacerda-Neto LJ, Figueiredo PRL, Oliveira LR, Rodrigues CKS, Sales VS, Figueiredo FRSDN, Nascimento EP, Monteiro AB, Amaro EN, Costa JGM, Coutinho HDM, de Menezes IRA, Kerntopf MR.. Antiulcerogenic Activity of the Hydroalcoholic Extract of Leaves of *Croton campestris* A. St.-Hill in Rodents. *Evid Based Complem Altern Med.* 2013; Article ID579346, 10 p.
- Critchfield JW, Butera ST, Folks TM. Inhibition of HIV activation in latently infected cells by flavonoid compounds. *AIDS Res Hum Retrovir.* 1996;12:39-46.
- Carotenuto A, Fattorusso E, Lanzotti V, Magno S, De Feo V, Cicala C. The flavonoids of *Allium neapolitanum*. *Phytochemistry.* 1997;44:949-57.
- Cheong H, Ryu SY, Oak MH, Cheon SH, Yoo GS, Kim KM. Studies of structure activity relationship for the anti-allergic actions. *Arch Pharmacol Res.* 1998;21:478-80.
- Costa AC. *Farmacognosia.* 2 ed. Lisboa: Fundação Calouste Gulbenkian; 1982. v.3
- Curtis HG, MacNaughton WK, Gall DG, Wallace JL. Intraluminal pH modulates gastric prostaglandin synthesis. *Can J Physiol Pharmacol.* 1995;73:130–4.

- Dekanski D, Ristic S, Mitrovic DM. Antioxidant effect of dry olive (*Olea europaea* L.) leaf extract on ethanol-induced gastric lesions in rats. *Mediterr J Nutr Metab*. 2009;2: 205–11.
- Ferreira O, Pinho SP. Solubility of flavonoids in pure solvents. *Ind Eng Chem Res*. 2012;51:6586–90.
- Giulietti AM, Harley RM, Queiroz LP, Wanderley MG, Van Den Berg C. Biodiversidade e conservação das plantas no Brasil. *Megadiversidade*. 2005;1(1):52-61.
- Golbabapour S, Hajrezaie M, Hassandarvish P, AbdulMajid N, Hadi AA, Nordin N, Abdulla MA. Acute Toxicity and Gastroprotective Role of *M. pruriens* Ethanol-Induced Gastric Mucosal Injuries in Rats. *BioMed Res Int*. 2013;13 pages.
- Hirano H, Takase H, Yamamoto K, Abe K, Saito J. Nobiletin as ulcer inhibitor. Patent-Japan KokaiTokkyoKoho.1994;06 72:870-874.
- Izzo AA, Di Carlo G, Mascolo N, Capasso F. Antiulcer effect of flavonoids: role of endogenous PAF. *Phytother Res*. 1994;8:179-181.
- Harborne JB. *Phytochemicals Methods*. London: Chapman and Hall; 1973. p. 49-188.
- Harborne JB. *Phytochemical Methods - A Guide to Modern Techniques of plant analysis*. London: Chapman and Hall; 1998.
- Havsteen BH. The biochemistry and medical significance of the flavonoids. *Pharmacol Ther*. 2002;96:67-202.
- Jamal A, Javed K, Aslam M, Jafri MA. Gastroprotective effect of cardamom, *Elettaria cardamomum* Maton. fruits in rats. *J Ethnopharmacol*. 2006;103(2):149–53.
- Lee B, Nugroho A, Bachri MS, Choi J, Lee KR, Choi JS, Kim WB, Lee KT, Lee JD, Park HJ. Anti-ulcerogenic Effect and HPLC Analysis of the Caffeoylquinic Acid-Rich Extract from *Ligularia stenocephala*. *Biol Pharm Bull*. 2010;33(3):493-7.
- Lorenzi H. *Plantas Mediciniais no Brasil*. Nova Odessa, (SP): Plantarum; 2002.
- Manach C, Morand C, Demigne C, Texier O, Regeat F, Révész C. Bioavailability of rutin and quercetin in rats. *FEBS Lett*. 1997;409:12-6.
- Martins RE, Castro DM, Castellani DC, Dias JE. *Plantas Mediciniais*. 1998; Ed. FV: Viçosa, p. 155.
- Matsumoto M, Hara H, Chiji H, Kasai K. Gastroprotective Effect of Red Pigments in Black Chokeberry Fruit (*Aronia melanocarpa* Elliot) on Acute Gastric Hemorrhagic Lesions in Rats. *J Agric Food Chem*. 2004;52:2226–9.
- Mizui T, Doteuchi M. Effect of polyamines on acidified ethanol induced gastric lesion in rats. *Jpn J Pharmacol*.1983;33:939-945.
- Morimoto Y, Shimohara K, Oshima S, Sukamoto T. Effects of the new anti-ulcer agent KB-5492 on experimental gastric mucosal lesions and gastric mucosal defensive factors, as compared to those of teprenone and cimetidine. *Jpn J Pharmacol*. 1991;57: 495–505.
- Mota KSL, Dias GEN, Pinto MEF, Ferreira AL, Souza-Brito ARM, Hiruma-Lima CA, Barbosa-Filho JM, Batista LM. Flavonoids with Gastroprotective Activity. *Molecules*. 2009;14:979-1012.
- Pepato MT, Keller EH, Baviera AM, Ketelhut IC, Vendramini RC, Brunetti IL. Anti-diabetic activity of *Bauhinia forficata* decoction in streptozotocin-diabetic rats. *J Ethnopharmacol*. 2002;81(2):191-7.
- Pires AM, Araújo PS. Risk perception and concepts about medicinal plants, herbal and allopathic medicines among pregnant women. *RBSP*. 2011;35(2):320–33.
- Pizzolatti MG, Cunha Jr A, Szpoganicz A, de Souza E, Braz-Filho R, Schripsena J. Flavonóides glicolisados das folhas verdes de *Bauhinia forficata*. *Quim Nova*. 2003; 26(4):466-9.
- Rainsford KD. The effect of 5-lipoxygenase inhibitors and the leukotriene antagonists on the development of gastric lesions induced by non-steroidal anti-inflammatory drugs in mice. *Agents and Actions*. 1978;21:316-9.
- Rong T. Chemistry and biochemistry of dietary polyphenols, Review. *Nutrients*. 2010;2(12):1231–46.
- Samonina GE, Kopylova GN, Lukjanzeva GV (2004). Antiulcer effects of amylin: a review. *Pathophysiol*. 2004;11(1):1–6.
- Szelenyi I, Thiemer K. Distention Ulcer as a Model for Testing of Drugs for Ulcerogenic Side-Effects. *Arch Toxicol*. 1978;41(1):99-105.
- Zuanazzi JAS, Montanha JA. Flavonóides. In: Simões CMO (org.) *Farmacognosia: da planta ao medicamento*. 6a. ed. Porto Alegre/Florianópolis: UFRGS/UFSC; 2010. p. 577-614.

Received on December 5th 2013

Accepted on February 27th 2014

